| lame: | |
|-------------|-------|
| APPT: Date: | Time: |

Instructions for Allergy Testing

We would like to review these important instructions regarding any medications you may be taking. Some medications interfere with the results – others may counteract your medications, resulting in an adverse effect.

Please do not take these medications **7 days** prior to your allergy test:

| Allegra (Fexofenadine) Allegra D | Tagamet |
|-------------------------------------|------------------------|
| Benadryl (Diphenhydramine) | Zantac |
| Atarax | Tylenol PM |
| Chlorpheniramine | Midol |
| Dramamine | Excedrin PM |
| Novahistine | Phenergan |
| Claritin (Loratadine) or Claritin D | Advil PM |
| Zyrtec | Xyzal (Levocetirizine) |

Avoid any over-the-counter preparations for cold, sinus or allergy. If in doubt, please call the office. If possible, inhalers should be avoided for **6 hours** before testing:

• Azmacort • Proventil • Ventolin • Flovent

If you take Beta Blockers please discuss this with us before your test or any allergy injections. If taking any of these medications you may need to hold your morning dose the day of your test.

Beta Blocker Medication List

| Brand Name | Generic Name |
|----------------------------------|----------------------------------|
| Bystolic | Nebivolol |
| Coreg (Carvedilol) | Carvedilol |
| Nadolol | Nadolol |
| Inderal/Inderal LA (Propranolol) | Propranolol |
| Lopressor | Metoprolol |
| Lopressor HCT | Metoprolol/HCTZ |
| Normodyne | Labetalol |
| Tenormin | Atenolol |
| Toprol XL | Metoprolol |
| Ziac | Bisoprolol & Hydrochlorothiazide |

The test will take 30-45 minutes. The test may be shortened and/or administered over multiple visits based on patient reactivity to the testing process. If it is necessary to cancel your test, please notify us at least 2 business days prior to you appointment. If any questions regarding any medications please contact clinic.

Please call (210)802-3777 if you have any questions.